

Racord® Plus

Twisted Gingival Retraction Cord with Epinephrine plus Astringent

Racord Plus is twisted cotton cord saturated with a solution of racemic epinephrine hydrochloride and astringent (zinc phenolsulfonate) and dried by a technique which retains the activity of the medication. Normal tissue moisture reactivates the medication, which promptly retracts gingival tissue in a manner that causes no tissue damage, yet fully exposes the gingival margin. The combination of epinephrine and astringent provides the cord with enhanced physical and chemical retraction.

Racord Plus is designed for dentists who insist on the best retraction medium available (epinephrine impregnated Racord), yet who are concerned with the cardiovascular effects of epinephrine which occur in some patients. By adding an astringent to the cord, the epinephrine content can be dropped by as much as 70% without sacrificing the effectiveness of full-strength epinephrine.

Contents: Each bottle contains 72 inches twisted cotton cord. The average concentration of medicament is:

	Epinephrine	Zinc Phenolsulfonate
Racord Plus 00	0.17 mg/in	0.30 mg/in
Racord Plus 0	0.225 mg/in	0.40 mg/in
Racord Plus 1	0.30 mg/in	0.70 mg/in
Racord Plus 2	0.35 mg/in	1.45 mg/in

Indications for Use: Use whenever gingival retraction is advisable in dental practice, as preceding impression taking with any elastic impression material; or prior to placement of caps, crowns, bridges; and during tooth preparation.

Contraindications: Racord Plus is contraindicated for persons who are epinephrine sensitive, or for persons suffering from cardiovascular disease, diabetes, or hyperthyroidism. Caution should also be taken for elderly, pregnant, and pediatric patients. For such persons Pascord® is the retraction cord of choice. It is packaged in four sizes, like Racord Plus, but contains no epinephrine.

Racord Plus contains only one third the epinephrine found in Racord and most other leading epinephrine-type retraction cords. Therefore the possibility of adverse epinephrine reactions is reduced accordingly. However, with the presence of epinephrine, the patient's medical history should be consulted and the usual precautions observed.

Warnings: Contains Sodium Bisulfite, a sulfite that may cause allergic-type reactions, including anaphylactic symptoms, and life-threatening or less severe asthmatic episodes, in certain susceptible persons. The overall prevalence of sulfite sensitivity in the general population is unknown, but probably low. Sulfite sensitivity is seen more in asthmatic than non-asthmatic persons.

**OPEN
HERE**

RX ONLY (In USA)

For professional use only. Read package insert for additional cautions and warnings. This product is not sterile. Single use only. Dispensed portions not to be reused.



**2929 Northrup Way
Bellevue, WA 98004 USA**



www.w.pascaldental.com



**2929 Northrup Way
Bellevue, WA 98004 USA
425.827.4694 800.426.8051
www.pascaldental.com**

visit us at pascaldental.com.

For more information about Pascal's cords and latest gingival retraction materials please

Racord Plus contains no artificial flavors.

Pascal's Racord Plus is free of gluten, soy, milk, egg, peanut, and tree nut products. Pascal directly.

For product SDS information please see company website at pascaldental.com or contact

immediately after use.

Dispensed portions of cord are intended for single use only and should be disposed of im-

then be removed from the mouth.

- Racord Plus completely retracts tissue within five to eight minutes and should suffice.

- Alternately, a single loop of cord will

be removed also.

- Once the impression procedure has been completed the smaller cord is re-

proceedure has been completed.

the gingival margin until the impression

leaving the smaller cord in place below

material, remove the larger cord loop only,

- When ready to apply the impression ma-

the smaller cord in the same manner.

- Apply a second loop of larger cord over

pack the loop firmly into the sulcus.

- Trim the ends for a slight overlap, and

tooth and over the crevice.

loop a strand of smaller cord around the

- For most effective use of Racord Plus,

Directions for Use:

mouth per appointment.

equivalent of one normal quadrant of the

an epinephrine cord should be limited to the

low), if sulcus space allows. Retraction with

may be packed around each tooth (see be-

be retracted, with a slight overlap. Two loops

the way around the neck of each tooth to

appropriate sizes as is necessary to go all

Amount to Use: Use as much cord of the

sulfonate allergy or sensitivity.

- Do not use on patients with zinc phenol-

allergy or sensitivity.

- Do not use on patients with epinephrine

Keep out of reach of children

- For topical intraoral use only. Do not

ingest.

- Read and follow all instructions.

Precautions: